

Gravitational Localization of Matters in 6D

Pavle Midodashvili* and Levan Midodashvili†

Department of Physics, Tskhinvali State University, 2 Besiki Str., Gori 383500, GEORGIA

(Dated: August 5, 2003)

We present a new 3-brane solution to Einstein's equations in (1+5)-spacetime with a negative bulk cosmological constant. This solution is a stringlike defect solution with decreasing scale function approaching a finite non-zero value in the radial infinity. It is shown that all local fields are localized on the brane only through the gravitational interaction.

It is believed that the idea of extra dimensions would be one of the most attractive ideas concerning unification of gauge fields with general relativity. Large extra dimensions offer an opportunity for a new solutions to old problems (smallness of cosmological constant, the origin of the hierarchy problem, the nature of flavor, etc.). This idea has been much investigated since the appearance of papers [1]. In such theories our world can be associated with a 3-brane, embedded in a higher-dimensional space-time with non-compact extra dimensions and non-factorizable geometry. In this scenario it is assumed that all the matter fields are constrained to live on the 3-brane and the corrections to four dimensional Newton's gravity low from bulk gravitons are small for macroscopic scales. But this models still need some natural mechanism of localization of known particles on the brane. The questions of matter localization on the brane has been investigated in various papers [2, 3, 4]. In our opinion the localizing force must be universal for all types of 4- dimensional matter fields. In our world the gravity is known to be the unique interaction which has universal coupling with all matter fields. So if extended extra dimensions exist, it is natural to assume that trapping of matter on the brane has a gravitational nature. It is of interest that recently in [6] it was found the model in the brane world where all the local bulk fields (ranging from the spin 0 scalar field to the spin 2 gravitational field) are localized on the 3-brane only by the universal interaction, i.e., the gravity. The solution is found in (1+5)-spacetime for the positive bulk cosmological constant $\Lambda > 0$ and has increasing scale factor $\phi(r)$ asymptotically approaching finite value at the radial infinity. Recently in [7] this solution was extended to the case of a general (p-1)-brane model with codimension n in general $D = p + n$ space-time dimensions.

In this article we introduce the new 3-brane solution in (1+5)-spacetime for the negative cosmological constant $\Lambda < 0$ which has decreasing scale factor $\phi(r)$ asymptotically approaching finite non-zero value at the radial infinity. We explicitly show that whole local fields (spin 0 scalar field, spin $1/2$ spinor field, spin 1 gauge field, spin $1/2$ gravitino field and spin 2 gravitational field) as well as totally antisymmetric tensor fields are localized on the 3-brane by the gravity.

Let us begin with the details of our solution. In 6D the Einstein equations with a bulk cosmological constant Λ and stress-energy tensor T_{AB}

$$R_{AB} - \frac{1}{2}g_{AB}R = \frac{1}{M^4}(\Lambda g_{AB} + T_{AB}) \quad (1)$$

can be derived from the whole action of the gravitating system

$$S = \int d^6x \sqrt{-g} \left[\frac{M^4}{2} (R + 2\Lambda) + L \right] \quad (2)$$

R_{AB} , R , M and L are respectively the Ricci tensor, the scalar curvature, the fundamental scale and the Lagrangian of matter fields (including brane). All of these physical quantities refer to (1+5)- space with signature $(+ - \dots -)$, capital Latin indices run over $A, B, \dots = 0, 1, 2, 3, 5, 6$. Suppose that the equations (1) admit a solution that is consistent with four-dimensional Poincaré invariance. Introducing for the extra dimensions the polar coordinates (r, θ) , where $0 \leq r < +\infty$, $0 \leq \theta < 2\pi$, the six-dimensional metric satisfying this ansatz we can choose in the form [4, 6]:

$$ds^2 = \phi^2(r) \eta_{\alpha\beta} (x^\nu) dx^\alpha dx^\beta - g(r) (dr^2 + r^2 d\theta^2), \quad (3)$$

where small Greek indices $\alpha, \beta, \dots = 0, 1, 2, 3$ numerate coordinates and physical quantities in four-dimensional space, the functions $\phi(r)$ and $g(r)$ depend only on r and are cylindrically symmetric in the extra-space, the metric signature is $(+ - \dots -)$. The function $g(r)$ must be positive to fix the signature of the metric (3).

*Electronic address: midodashvili@hotmail.com

†Electronic address: Levmid@hotmail.com

The source of the brane is described by a stress-energy tensor T_{AB} also cylindrically symmetric in the extra-space. Its nonzero components we choose in the form

$$T_{\alpha\beta} = -g_{\alpha\beta}F_0(r), \quad T_{ij} = -g_{ij}F(r), \quad (4)$$

where we have introduced two source functions F_0 and F , which depend only on the radial coordinate r .

By using cylindrically symmetric metric ansatz (3) and stress-energy tensor (4), the Einstein equations become

$$3\frac{\phi''}{\phi} + 3\frac{\phi'^2}{\phi^2} + 3\frac{\phi'}{r\phi} + \frac{g''}{2g} - \frac{g'^2}{2g^2} + \frac{g'}{2rg} = \frac{g}{M^4}(F_0 - \Lambda) + \frac{g}{\phi^2} \frac{\Lambda_{phys}}{M_P^2}, \quad (5)$$

$$6\frac{\phi'^2}{\phi^2} + 2\frac{g'\phi'}{g\phi} + 4\frac{\phi'}{r\phi} = \frac{g}{M^4}(F - \Lambda) + 2\frac{g}{\phi^2} \frac{\Lambda_{phys}}{M_P^2}, \quad (6)$$

$$4\frac{\phi''}{\phi} + 6\frac{\phi'^2}{\phi^2} - 2\frac{g'\phi'}{g\phi} = \frac{g}{M^4}(F - \Lambda) + 2\frac{g}{\phi^2} \frac{\Lambda_{phys}}{M_P^2}, \quad (7)$$

where the prime denotes differentiation d/dr . The constant Λ_{phys} represents the physical four-dimensional cosmological constant, where

$$R_{\alpha\beta}^{(4)} - \frac{1}{2}g_{\alpha\beta}R^{(4)} = \frac{\Lambda_{phys}}{M_P^2}g_{\alpha\beta}. \quad (8)$$

In this equation $R_{\alpha\beta}^{(4)}$, $R^{(4)}$ and M_P are four-dimensional physical quantities: Ricci tensor, scalar curvature and Planck scale.

In the case $\Lambda_{phys} = 0$ from the equations (5), (6) and (7) we can find

$$F' + 4\frac{\phi'}{\phi}(F - F_0) = 0, \quad (9)$$

$$g = \frac{\delta\phi'}{r}, \quad \delta = const, \quad (10)$$

$$r\frac{\phi''}{\phi} + 3r\frac{\phi'^2}{\phi^2} + \frac{\phi'}{\phi} = \frac{rg}{2M^4}(F - \Lambda), \quad (11)$$

where δ denotes the integration constant. The (9) represents the connection between source functions, it is simply a consequence of the conservation of the stress-energy tensor and can be also independently derived directly from $D_A T_B^A = 0$.

Suppose that the "width" of brane sitting in the origin $r = 0$ is equal to ε . It is worthwhile to mention that the brane is assumed to have the nonvanishing "brane width" since the "brane width" ε appears in the later arguments of localization of the bulk fields and plays a role as a short-distance cutoff [7]. Outside a core of radius ε we take the source functions in the form $F(r) = f\phi^{-2}$, where f is some constant. Taking the first integral of the last equation [3], we get

$$r\phi' = -\frac{\delta\Lambda}{10M^4}\left(\phi^2 - \frac{5f}{3\Lambda}\right) + \frac{C}{\phi^3}, \quad (12)$$

where C is the integration constant. Setting $C = 0$, introducing the positive parameters

$$a = \frac{\delta\Lambda}{10M^4} > 0, \quad d^2 = \frac{5f}{3\Lambda} > 0, \quad (13)$$

and imposing boundary condition at the infinity of transverse space

$$\phi(r)|_{r=+\infty} = const > 0, \quad (14)$$

we can easily find two solutions of the equation (12) in the following cases:

i) $\Lambda > 0, f > 0, \delta > 0$

$$\phi(r) = d \frac{r^b - c}{r^b + c}, \quad g(r) = 2\delta db \frac{cr^{b-2}}{(r^b + c)^2}; \quad (15)$$

ii) $\Lambda < 0, f < 0, \delta < 0$

$$\phi(r) = d \frac{r^b + c}{r^b - c}, \quad g(r) = 2|\delta| db \frac{cr^{b-2}}{(r^b - c)^2}; \quad (16)$$

where $c > 0$ is positive integration constant and we have introduced the parameter $b = 2ad$. The first solution (15) is the same one found in [6]. This solution exists in the case of positive bulk cosmological constant $\Lambda > 0$. The scale factor $\phi(r)$ of this solution is increasing function asymptotically approaching the finite value at the radial infinity. The second solution (16) is the new one. It exists for the negative cosmological constant $\Lambda < 0$, and its scale factor $\phi(r)$ is decreasing function asymptotically approaching the finite non-zero value at the radial infinity. In this article we consider the second solution (the first one has been already examined in [6]).

To avoid singularities on the brane (sitting at the origin $r = 0$) we take the boundary conditions for the solution in the form

$$\phi(r)|_{r=\varepsilon} = 1, \quad \phi(r)|_{r=+\infty} = d. \quad (17)$$

In (17) ε denotes the "brane width", which takes a fixed value. This boundary conditions allow us to express the integration constant c in terms of the "brane width":

$$c = \frac{1-d}{1+d} \varepsilon^b, \quad \text{where } 0 < d < 1. \quad (18)$$

Now we turn our attention to the problem of the localization of the bulk fields on the brane in the background geometry (16). Of course, in due analysis, we will neglect the back-reaction on the geometry induced by the existence of the bulk fields, and from now on, without loss of generality, we shall take a flat metric on the brane.

We start with a massless, spin 0, real scalar coupled to gravity:

$$S_0 = -\frac{1}{2} \int d^6x \sqrt{-6} g g^{AB} \partial_A \Phi \partial_B \Phi. \quad (19)$$

The corresponding equation of motion has the form

$$\frac{1}{\sqrt{-6}g} \partial_M \left(\sqrt{-6}g g^{MN} \partial_N \Phi \right) = 0. \quad (20)$$

It turns out that in the background metric (16) the zero-mode solution of (20) is $\Phi_0(x^M) = v(x^\mu) \rho_0$, where $\rho_0 = \text{constant}$, and $v(x^\mu)$ satisfies the Klein-Gordon equation on the brane $\eta^{\mu\nu} \partial_\mu \partial_\nu v(x^\mu) = 0$. Substituting this solution into the starting action (19), the action can be cast to

$$S_0 = -\frac{1}{2} \rho_0^2 \int_0^{2\pi} d\theta \int_\varepsilon^{+\infty} dr \phi^2 g r \int d^4x \sqrt{-\eta} \eta^{\mu\nu} \partial_\mu v(x^\alpha) \partial_\nu v(x^\alpha) + \dots = \quad (21)$$

$$= -\pi \delta \rho_0^2 \int_{\phi(\varepsilon)}^{\phi(+\infty)} \phi^2 d\phi \int d^4x \sqrt{-\eta} \eta^{\mu\nu} \partial_\mu v(x^\alpha) \partial_\nu v(x^\alpha) + \dots = \quad (22)$$

$$= -\frac{\pi |\delta| \rho_0^2}{3} (1 - d^3) \int d^4x \sqrt{-\eta} \eta^{\mu\nu} \partial_\mu v(x^\alpha) \partial_\nu v(x^\alpha) + \dots \quad (23)$$

The integral over r in (21) is finite, so the 4-dimensional scalar field is localized on the brane.

For spin $1/2$ fermion starting action is the Dirac action given by

$$S_{\frac{1}{2}} = \int d^6x \sqrt{-6} g \bar{\Psi} i \Gamma^M D_M \Psi, \quad (24)$$

from which the equation of motion is given by

$$\Gamma^M D_M \Psi = (\Gamma^\mu D_\mu + \Gamma^r D_r + \Gamma^\theta D_\theta) \Psi = 0. \quad (25)$$

We introduce the vielbein $h_A^{\tilde{A}}$ through the usual definition $g_{AB} = h_A^{\tilde{A}} h_B^{\tilde{B}} \eta_{\tilde{A}\tilde{B}}$ where $\tilde{A}, \tilde{B}, \dots$ denote the local Lorentz indices. Γ^A in a curved space-time is related to γ^A by $\Gamma^A = h_A^{\tilde{A}} \gamma^{\tilde{A}}$. The spin connection $\omega_M^{\tilde{M}\tilde{N}}$ in the covariant derivative $D_M \Psi = \left(\partial_M + \frac{1}{4} \omega_M^{\tilde{M}\tilde{N}} \gamma_{\tilde{M}\tilde{N}} \right) \Psi$ is defined as

$$\omega_M^{\tilde{M}\tilde{N}} = \frac{1}{2} h^{N\tilde{M}} \left(\partial_M h_N^{\tilde{N}} - \partial_N h_M^{\tilde{N}} \right) - \frac{1}{2} h^{N\tilde{N}} \left(\partial_M h_N^{\tilde{M}} - \partial_N h_M^{\tilde{M}} \right) - \frac{1}{2} h^{P\tilde{M}} h^{Q\tilde{N}} \left(\partial_P h_{Q\tilde{R}} - \partial_Q h_{P\tilde{R}} \right) h_M^{\tilde{R}}.$$

After these conventions are set we can decompose the 6-dimensional spinor into the form $\Psi(x^M) = \psi(x^\mu) A(r) \sum e^{il\theta}$. We require that the four-dimensional part satisfies the massless equation of motion $\gamma^\mu \partial_\mu \psi(x^\beta) = 0$ and the chiral condition $\gamma^r \psi(x^\mu) = \psi(x^\mu)$. As a result we obtain the following equation for the s -wave mode

$$\left[\partial_r + 2 \frac{\phi'}{\phi} + \frac{1}{2} \frac{\partial_r (r g^{\frac{1}{2}})}{r g^{\frac{1}{2}}} \right] A(r) = 0. \quad (26)$$

The solution to this equation reads:

$$A(r) = A_0 \phi^{-2} g^{-\frac{1}{4}} r^{-\frac{1}{2}}, \quad (27)$$

with A_0 being an integration constant. Substituting this solution into the Dirac action (24) we have

$$S_{\frac{1}{2}} = 2\pi A_0^2 \int_\varepsilon^{+\infty} dr \phi^{-1} g^{\frac{1}{2}} \int d^4 x \sqrt{-\eta} \bar{\psi} i \gamma^\mu \partial_\mu \psi + \dots = \quad (28)$$

$$= 2\pi A_0^2 \sqrt{\frac{2|\delta|bc}{d}} \int_\varepsilon^{+\infty} \frac{r^{\frac{1}{2}b} dr}{r(r^b + c)} \int d^4 x \sqrt{-\eta} \bar{\psi} i \gamma^\mu \partial_\mu \psi + \dots. \quad (29)$$

As long as $b > 0$ and the "brane width" ε is nonvanishing the integral over r is obviously finite. Indeed, the integrand in (29) scales as $r^{-\frac{1}{2}b-1}$ at the radial infinity and is the smooth functions between $r = \varepsilon$ and $r = +\infty$, so this integral over r is finite even if the analytic expression is not available. So the spin $1/2$ fermion is localized on the brane only by the gravitational interaction.

Now let us consider the action of $U(1)$ vector field:

$$S_1 = -\frac{1}{4} \int d^6 x \sqrt{-6g} g^{AB} g^{MN} F_{AM} F_{BN}, \quad (30)$$

where $F_{MN} = \partial_M A_N - \partial_N A_M$ as usual. From this action the equation of motion is given by

$$\frac{1}{\sqrt{-6g}} \partial_M \left(\sqrt{-6g} g^{MN} g^{RS} F_{NS} \right) = 0. \quad (31)$$

By choosing the gauge condition $A_\theta = 0$ and decomposing the vector field as

$$A_\mu(x^M) = a_\mu(x^\mu) \sum_{l,m} \sigma_m(r) e^{il\theta}, \quad (32)$$

$$A_r(x^M) = a_r(x^\mu) \sum_{l,m} \sigma_m(r) e^{il\theta}, \quad (33)$$

it is straightforward to see that there is the s -wave ($l = 0$) constant solution $\sigma_m(r) = \sigma_0 = \text{const}$ and $a_r = \text{const}$. In deriving this solution we have used $\partial_\mu a^\mu = \partial^\mu f_{\mu\nu} = 0$ with the definition of $f_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu$. As in the previous cases, let us substitute this constant solution into the action (30). It turns out that the action is reduced to

$$S_1 = -\frac{\pi}{2} \sigma_0^2 \int_\varepsilon^{+\infty} dr g r \int d^4 x \sqrt{-\eta} \eta^{\alpha\beta} \eta^{\mu\nu} f_{\alpha\mu} f_{\beta\nu} + \dots = \quad (34)$$

$$= \frac{\pi |\delta|}{2} \sigma_0^2 \int_{\phi(\varepsilon)}^{\phi(+\infty)} d\phi \int d^4x \sqrt{-\eta} \eta^{\alpha\beta} \eta^{\mu\nu} f_{\alpha\mu} f_{\beta\nu} + \dots = -\frac{\pi |\delta|}{2} \sigma_0^2 (1-d) \int d^4x \sqrt{-\eta} \eta^{\alpha\beta} \eta^{\mu\nu} f_{\alpha\mu} f_{\beta\nu} + \dots \quad (35)$$

As we can see from (35) the integral over r in (34) is finite. Thus, the vector field is localized on the brane. Next we consider spin $3/2$ field (the gravitino). We begin with the action of the Rarita-Schwinger gravitino

$$S_{\frac{3}{2}} = \int d^6x \sqrt{-6g} \bar{\Psi}_A i \Gamma^{[A} \Gamma^B \Gamma^{C]} D_B \Psi_C, \quad (36)$$

from which the equation of motion is given by

$$\Gamma^{[A} \Gamma^B \Gamma^{C]} D_B \Psi_C = 0. \quad (37)$$

Here the square bracket denotes the anti-symmetrization and the covariant derivative is defined with the affine connection $\Gamma_{BC}^A = h_{\tilde{B}}^A \left(\partial_B h_{\tilde{C}}^{\tilde{B}} + \omega_{\tilde{B}}^{\tilde{B}\tilde{C}} h_{C\tilde{C}} \right)$ by

$$D_A \Psi_B = \partial_A \Psi_B - \Gamma_{AB}^C \Psi_C + \frac{1}{4} \omega_{\tilde{A}}^{\tilde{A}\tilde{B}} \gamma_{\tilde{A}\tilde{B}} \Psi_B \quad (38)$$

After taking the gauge condition $\Psi_\theta = 0$ we look for the solutions of the form $\Psi_\mu(x^A) = \psi_\mu(x^\nu) u(r) \sum e^{i l \theta}$, $\Psi_r(x^A) = \psi_r(x^\nu) u(r) \sum e^{i l \theta}$ where $\psi_\mu(x^\nu)$ satisfies the following equations $\gamma^\nu \psi_\nu = \partial^\mu \psi_\mu = \gamma^{[\nu} \gamma^\rho \gamma^{\tau]} \partial_\rho \psi_\tau = 0$, $\gamma^r \psi_\nu = \psi_\nu$. For the s -wave solution and $\psi_r(x^\nu) = 0$ we have for the equation of motion (37) the following form

$$\left[\partial_r + \frac{3}{2} \frac{\phi'}{\phi} + \frac{1}{2} \frac{\partial_r (r g^{\frac{1}{2}})}{r g^{\frac{1}{2}}} \right] u(r) = 0. \quad (39)$$

The solution to this equation is

$$u(r) = u_0 \phi^{-\frac{3}{2}} g^{-\frac{1}{4}} r^{-\frac{1}{2}}, \quad (40)$$

with u_0 being an integration constant. Substituting this solution into the action (36) we get

$$S_{\frac{3}{2}} = 2\pi u_0^2 \int_{\varepsilon}^{+\infty} dr \phi^{-2} g^{\frac{1}{2}} \int d^4x \sqrt{-\eta} \bar{\psi}_\mu i \gamma^{[\mu} \gamma^\nu \gamma^{\rho]} \partial_\nu \psi_\rho + \dots = \quad (41)$$

$$= 2\pi u_0^2 \sqrt{\frac{2|\delta| b c_2}{d}} \int_{\varepsilon}^{+\infty} \frac{r^{\frac{1}{2}b} (r^b - c_2)}{r (r^b + c_2)^2} dr \int d^4x \sqrt{-\eta} \bar{\psi}_\mu i \gamma^{[\mu} \gamma^\nu \gamma^{\rho]} \partial_\nu \psi_\rho + \dots \quad (42)$$

As in the case of spin $1/2$ fermion (29) the integrand in (42) scales as $r^{-\frac{1}{2}b-1}$ at the radial infinity and is smooth function between $r = \varepsilon$ and $r = +\infty$, so the integral over r is finite as long as the "brane width" ε is non-zero. This means that spin $3/2$ field is localized on the brane.

Now let us consider spin 2 gravitational field. In this case we consider the spin-2 metric fluctuations $H_{\mu\nu}$:

$$ds^2 = \{ \phi^2(r) \eta_{\alpha\beta}(x^\nu) + H_{\alpha\beta} \} dx^\alpha dx^\beta - g(r) (dr^2 + r^2 d\theta^2). \quad (43)$$

The corresponding equation of motion for the fluctuations has the following form:

$$\frac{1}{\sqrt{-6g}} \partial_A \left(\sqrt{-6g} g^{AB} \partial_B H_{\mu\nu} \right) = 0. \quad (44)$$

From the 4-dimensional point of view fluctuations are described by a tensor field $H_{\mu\nu}$ which is transverse and traceless:

$$\partial_\mu H_\nu^\mu = 0, \quad H_\mu^\mu = 0. \quad (45)$$

This tensor is invariant under 4-dimensional general coordinate transformations. We look for solutions of the form

$$H_{\alpha\beta} = h_{\alpha\beta}(x^\nu) \sum_{ml} \tau_m(r) \exp(i\theta l), \quad (46)$$

where $\partial^2 h_{\mu\nu}(x^\alpha) = m_0^2 h_{\mu\nu}(x^\alpha)$. It is easy to show that the equation of motion (44) has the zero-mass ($m_0 = 0$) and s -wave ($l = 0$) constant solution $\tau_0 = \text{const}$. Substitution of this zero mode into the Einstein-Hilbert action leads to

$$S_2 \sim 2\pi\tau_0^2 \int_\varepsilon^{+\infty} dr \phi^2 gr \int d^4x [\partial^\rho h^{\alpha\beta} \partial_\rho h_{\alpha\beta} + \dots] = \quad (47)$$

$$= 2\pi\delta\tau_0^2 \int_{\phi(\varepsilon)}^{\phi(+\infty)} \phi^2 d\phi \int d^4x [\partial^\rho h^{\alpha\beta} \partial_\rho h_{\alpha\beta} + \dots] = \frac{2}{3}\pi |\delta| \tau_0^2 (1 - d^3) \int d^4x [\partial^\rho h^{\alpha\beta} \partial_\rho h_{\alpha\beta} + \dots]. \quad (48)$$

The integral over r is finite, so the bulk graviton is trapped on the brane. With our background metric the general expression for the four-dimensional Planck scale M_P , expressed in terms of M , is

$$M_P^2 = 2\pi M^4 \int_\varepsilon^{+\infty} dr \phi^2 gr = 2\pi\delta M^4 \int_\varepsilon^{+\infty} dr \phi^2 \phi' = \frac{2}{3}\pi |\delta| M^4 (1 - d^3). \quad (49)$$

The inequality $M \ll M_P$ is possible by adjusting M^4 and the product $|\delta|(1 - d^3)$, and thus could lead to a solution of the gauge hierarchy problem.

Finally, let us consider the totally antisymmetric tensor fields. The action of k -rank totally antisymmetric tensor field A_k is of the form

$$S_k = -\frac{1}{2} \int F_{k+1} \wedge *F_{k+1}, \quad (50)$$

where $F_{k+1} = dA_k$. The corresponding equation of motion is given by

$$d \wedge *F_{k+1} = 0. \quad (51)$$

It is easy to show that $A_{\mu_1\mu_2\dots\mu_k} = a_{\mu_1\mu_2\dots\mu_k}(x^\nu)u_0$ with $u_0 = \text{const}$ is a solution to the equation of motion (51) if $d \wedge *f = 0$ where $f = da$. Substituting this solution in the action (50) leads to the expression

$$S_k \sim \int_\varepsilon^{+\infty} dr \phi^{2-2k} gr \int f_{k+1} \wedge *f_{k+1} + \dots = \quad (52)$$

$$= \delta \int_{\phi(\varepsilon)}^{\phi(+\infty)} \phi^{2-2k} d\phi \int f_{k+1} \wedge *f_{k+1} + \dots = \frac{|\delta|}{(3-2k)} (1 - d^{3-2k}) \int f_{k+1} \wedge *f_{k+1} + \dots \quad (53)$$

As we can see from (53) the integral over r in (52) is finite, so the totally antisymmetric tensor fields are also localized on the brane by the gravitational interaction.

In conclusion, in this article we have presented a new 3-brane solution with a decreasing scale factor $\phi(r)$. This solution is found for the negative bulk cosmological constant $\Lambda < 0$. In addition we have presented a complete analysis of localization of a bulk fields on a brane via the gravitational interaction without including an additional interaction. The technical reason of the localization of all bulk fields lies not only in the fact that the scale factor $\phi(r)$ is a smooth function without singularities from the edge of the brane to the radial infinity and approaches a definite non-zero value at the infinity. The main reason is connected with the properties of the function $g(r)$. On the edge of the brane ($r = \varepsilon$) it has a value $g(\varepsilon) = a\delta(1 - d^2)\varepsilon^{-2}$ and fast enough tends to zero as one moves of the brane. In the case $\varepsilon \ll 1$, $\delta \gg 1$ and $b = 2ad = \frac{\delta\Lambda}{5M^4}d \gg 1$ this function has a δ -like behavior, and in contrast to the model considered in [7] in our model the wave functions of the zero-mode solutions of the bulk fields along the extra dimensions are peaked at the location of the brane.

Acknowledgements: P.M. would like to acknowledge the hospitality extended during his visits at the Abdus Salam International Centre for Theoretical Physics where the main part of this work was done.

-
- [1] V.A. Rubakov and M.E. Shaposhnikov, Phys. Lett. **B125** (1983) 136; Phys. Lett. **B125** (1983) 139.
 K. Akama, in *Lecture Notes in Physics, 176, Gauge Theory and Gravitation, Proceedings, Nara, Japan, 1982*,
 edited by K.Kikkawa, N.Nakanishi and H.Nariai, (Springer-Verlag, 1983) 267 [hep-th/0001113].
 M. Visser, Phys. Lett. **B159** (1985) 22.

- G.W. Gibbons and D.L. Wiltshire, Nucl. Phys. B287 (1987) 717 [hep-th/0109093].
- I. Antoniadis, Phys. Lett. **B246** (1990) 377.
- I. Antoniadis, N.Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B436**, (1998) 257.
- L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, (1999) 3370.
- N. Arkani-Hamed, S. Dimopoulos and G.Dvali, Phys. Lett. **B429** (1998) 263 ; Phys. Rev. **D59**, (1999) 086004.
- M. Gogberashvili, Phys. Lett. **B553**, (2003) 284; J. Math. Phys. **43**, (2002) 4886; J. Mod. Phys. **D11**, (2002) 1639.
- [2] Z. Chacko and A.E. Nelson, Phys. Rev. **D62** (2000) 085006.
- I. Oda, Phys. Rev. **D62** (2000) 126009.
- T. Gherghetta, E. Roessl and M. Shaposhnikov, Phys. Lett. **B491** (2000) 353.
- M. Giovannini, H. Meyer and M. Shaposhnikov, Nucl. Phys. **B619**, (2001) 615.
- Panagiota Kanti, Richard Madden and Keith A. Olive, Phys.Rev. **D64**, (2001) 044021 [hep-th/0104177].
- M. Giovannini, Phys. Rev. **D66**, (2002) 044016.
- S. Randjbar-Daemi and M. Shaposhnikov, Nucl. Phys. **B645**, (2002) 188.
- T. Gherghetta and M. Shaposhnikov, Phys. Rev. Lett. **85**, (2000) 240.
- A. Chodos and E. Poppitz, Phys. Lett. **B471**, (1999) 119.
- A.G. Cohen and D.B. Kaplan, Phys. Lett. **B470**, (1999) 52.
- B.Bajc and G. Gabadadze, Phys. Lett. **B474**, (2000) 282.
- [3] M. Gogberashvili and P. Midodashvili, Phys. Lett. **B515** (2001) 447.
- [4] M. Gogberashvili and P.Midodashvili, Europhys. Lett. **61** (2003) 308.
- [5] P. Midodashvili, hep-th/0308003.
- [6] M. Gogberashvili and D. Singleton, hep-th/0305241.
- [7] I. Oda, hep-th/0307119.